

Introduction

- Alveolar Echinococcosis (AE) is a rare zoonotic disease caused by the helminthic parasite *Echinococcus multilocularis*.^{1,2}
- E. multilocularis* is endemic to the northern hemisphere affecting approximately 0.03-1.2/100,000 individuals annually.^{1,2,3}
- Although more prevalent in Europe and Asia, the rates of AE have increased in North America in recent years.⁴
- AE is marked by visceral hydatid cyst formation, most commonly in the liver, with expansive, tumor-like growth and invasion.^{1,2,5}
- Similar to malignant neoplasms, AE infections are able to spread to adjacent or distant sites through direct, lymphatic or hematogenous pathways.^{1,3}
- Usually asymptomatic for a 5-15 year period and most commonly presents with abdominal pain, fatigue and weight loss.^{3,5,6}
- Often misdiagnosed as primary liver malignancies due to their tumor-like growth pattern.^{2,3,5}
- If inadequately treated or left untreated AE has a >90% fatality rate.^{5,7}

Clinical History

- A patient in their 30s presented with right upper quadrant pain, worsened after eating, along with concurrent shoulder pain.
- Ultrasound, CT and MRI scans revealed an ill-defined heterogeneous mass (10.5x 10.5 x15 cm) in the right lobe of the liver consistent with possible hepatocellular carcinoma.
- A right hepatectomy was performed to remove the mass along with portal and retroperitoneal lymph nodes.

Results

Gross Examination

- Sectioning revealed a light tan, ill-defined, variegated mass (17.1 x 10.2 x 9.7 cm) associated with a light tan nodular area of liver capsule.
- Inconsistent with the typical presentation of hepatocellular carcinoma, the mass consisted of tan to yellow solid components and mucinous to milky cystic components.
 - Hepatocellular carcinoma usually presents as a nodular (unifocal or multifocal) or diffusely infiltrative mass often arising in the background of cirrhosis.⁸
- The mass grossly abuts the liver capsule, resection margin and hepatic tubular structures.
- The remaining liver parenchyma had a nutmeg appearance.
- Representative sections of the mass and potential lymph nodes (0.7 and 2.7 cm in maximum dimension) were submitted for histological examination.




Fig. 1. Gross sections of the hepatic mass. (A) Solid and multicystic components on the cut surface of the mass with nutmeg appearance of uninvolved liver parenchyma. (B) Magnified view of solid and cystic hepatic mass.

Histology & Ancillary Studies

- Microscopic examination of the liver mass revealed the presence of laminated membranes and scolices consistent with echinococcal infections.
- A portion of fibroadipose tissue was also involved by echinococcosis.
- A final diagnosis of AE was determined after serology and microbiology were performed and confirmed an infection with *E. multilocularis*.




Fig. 2. Histological section of liver mass demonstrating *E. multilocularis*. (H&E, 40x magnification)




Fig. 3. Histological section demonstrating cystic spaces in liver. (H&E, 10x magnification)

Discussion

- Staging and classification of AE is based on the European Network for Concerted Surveillance of Alveolar Echinococcosis and the WHO Informal Group on Echinococcosis PNM system.^{6,9}

Table 1: PNM classification of alveolar echinococcosis.^{6,9}

P	Hepatic localization of the parasite
Px	Primary tumor cannot be assessed
P0	No detectable tumor in liver
P1	Peripheral lesion without proximal vascular and/or biliary involvement
P2	Central lesions with proximal vascular and/or biliary involvement of one lobe
P3	Central lesions with hilar or biliary involvement of both lobes and/or with involvement of two hepatic veins
P4	Any liver lesion with extension along the vessels (inferior vena cava, portal veins and arteries) and the biliary tree
N	Extra-hepatic involvement of neighbouring organs and regional lymph nodes
Nx	Not evaluable
N0	No regional involvement
N1	Involvement of neighbouring organs or tissues
M	Absence or presence of distant metastasis
Mx	Not completely assessed
M0	No metastasis on chest radiograph and computer tomography
M1	Metastasis present

Table 2: PNM stage grouping of alveolar echinococcosis.^{6,9}

Stage	Criteria		
I	P1	N0	M0
	P2	N0	M0
IIa	P3	N0	M0
	P1-3	N1	M0
IIb	P4	N0	M0
	P4	N1	M0
IV	Any P	Any N	M1

- Accurate descriptions of mass location, invasion to adjacent structures and lymph node involvement are all important for determining AE stage which directly impacts patient treatment.
- Treatment for AE may include antiparasitic drugs therapy, surgical resections, liver transplantation or combinations of the three.⁶

Conclusion

- AE is a rare zoonotic disease associated with a poor prognosis if left untreated.
- The characteristic tumor-like growth and dissemination patterns makes proper initial diagnosis difficult in areas with low prevalence.
- Careful grossing on echinococcosis cases by pathologists' assistants is of utmost importance for accurate staging and subsequent treatment.

References

- Buttenschoen K, Kern P, Reuter S, Barth TFE. Hepatic infestation of echinococcus multilocularis with extension to regional lymph nodes. *Langenbeck's Arch Surg*. 2009. doi:10.1007/s00423-009-0481-0
- Numari G, Pinzone MR, Gruttaduria S, et al. Hepatic echinococcosis: Clinical and therapeutic aspects. *World J Gastroenterol*. 2012. doi:10.3748/wjg.v18.i13.1448
- Chouhan MD, Wiley E, Chiodini PL, Amin Z. Hepatic alveolar hydatid disease (Echinococcus multilocularis), a mimic of liver malignancy: a review for the radiologist in non-endemic areas. *Clin Radiol*. 2019. doi:10.1016/j.crad.2019.01.007
- Massolo A, Klein C, Kowalewska-Grochowska K, et al. European Echinococcus multilocularis Identified in Patients in Canada. *N Engl J Med*. 2019. doi:10.1056/NEJMc1814975
- Pohnan R, Ryska M, Hytych V, Matej R, Hrabal P, Pudil J. Echinococcosis mimicking liver malignancy: A case report. *Int J Surg Case Rep*. 2017. doi:10.1016/j.ijscr.2017.04.032
- Mihmanli M, Idiz UO, Kaya C, et al. Current status of diagnosis and treatment of hepatic echinococcosis. *World J Hepatol*. 2016. doi:10.4254/wjh.v8.i28.1169
- Liccioli S, Duignan PJ, Lejeune M, Deunk J, Majid S, Massolo A. A new intermediate host for Echinococcus multilocularis: The southern red-backed vole (Myodes gapperi) in urban landscape in Calgary, Canada. *Parasitol Int*. 2013. doi:10.1016/j.parint.2013.03.007
- Schlageter M, Terracciano LM, D'Angelo S, Sorrentino P. Histopathology of hepatocellular carcinoma. *World J Gastroenterol*. 2014. doi:10.3748/wjg.v20.i43.15955
- Kern P, Wen H, Sato N, et al. WHO classification of alveolar echinococcosis: Principles and application. In: *Parasitology International*. 2006. doi:10.1016/j.parint.2005.11.041